




1/12



pLNH31, pLNH32, pLNH33, or pLNH 34

Fig. 1






Fermentation Time (hr)


Fermentation of Glucose and Xylose by LNH32

Fig. 2




Simultaneous Fermentation of Glucose and Xylose by Recombinant Saccharomyces LNH33

Fig. 3




Fermentation of Glucose and Xylose by the Un-Engineered Parent 1400 Saccharomyces Yeast

Fig. 4




Fermentation of glucose and xylose by LNH32 after being cultured for 4 and 20 generarions in non-selective (glucose) medium.

 $Fig.\,\,5$ SUBSTITUTE SHEET (RULE 20)




Fermentation of glucose and xylose by LNH33 after being cultured for 4 and 20 generations in non-selective (glucose) medium.

Fig. 6
SUBSTITUTE SHEET (RULE 26)




Fermentation of glucose and xylose by LNH-ST(1) after being cultured for 4, 20, and 40 generations in non-selective (glucose) medium.

Fig. 7
SUBSTITUTE SHEET (RULE 28)



- (A) Yeast (S. <u>cerevisiae</u>) AH22 cultured in YEPD (1% yeast extracts, 2% peptone, 2% glucose or YEP (1% yeast extracts, 2% peptone).
- (B) Yeast AH22 cultured in YEPXlu (1% yeast extracts, 2% peptone, 2% xylulose) or YEP.







Fig. 9b Substitute sheet (Rule 20)






Construction of pKS(-)-KK-AR-KD plasmids


†The XhoI site was regenerated after ligation; *Intact ADC1 promoter; ** ADC1 promoter with TRP5 ribosomal binding site

the start that the st



Recombinant Saccharomyces 1400(LNH-ST) for fermenting Glucose and Xylose

 $Fig. \ 11$ SUBSTITUTE SHEET (RULE 28)



12/12




Fig. 12